

The debate has gone back and forth over the last 10 years as new data are found, but measurements by Deep Impact/EPOXI, Cassini and most recently the Lunar Reconnaissance Orbiter and Chandrayaan-1 are now considered conclusive. Beneath the shadows of polar craters, and clinging to the lunar regolith, billions of gallons of water are available for harvesting by future astronauts.

The image to the left created by the Moon Minerology Mapper (M3) instrument onboard Chandrayaan-1, shows deposits and sources of hydroxyl molecules. The data has been colored blue and superimposed on a lunar photo.

Complimentary data from the Deep Impact/EPOXI and Cassini missions of the rest of the lunar surface also detected hydroxyl molecules covering about 25% of the surveyed lunar surface. The hydroxyl molecule consists of one atom of oxygen and one of hydrogen, and because water is basically a hydroxyl molecule with a second hydrogen atom added, detecting hydroxyl on the moon is an indication that water molecules are also present.

How much water might be present? The M3 instrument can only detect hydroxyl molecules if they are in the top 1-millimeter of the lunar surface. The measurements also suggest that about 1 metric ton of lunar surface has to be processed to extract 1 liter (0.26 gallons) of water.

Problem 1 – The radius of the moon is 1,731 kilometers. How many cubic meters of surface volume is present in a layer that is 1 millimeter thick?

Problem 2 – The density of the lunar surface (called the regolith) is about 3000 kilograms/meter³. How many metric tons of regolith are found in the surface volume calculated in Problem 1?

Problem 3 – The concentration of water is 1 liter per metric ton. How many liters of water could be recovered from the 1 millimeter thick surface layer if 25% of the lunar surface contains water?

Problem 4 – How many gallons could be recovered if the entire surface layer were mined? (1 Gallon = 3.78 liters).

Problem 1 – The radius of the moon is 1,731 kilometers. How many cubic meters of surface volume is present in a layer that is 1 millimeter thick?

Answer: The surface area of a sphere is given by $S = 4\pi r^2$ and so the volume of a layer with a thickness of L is $V = 4\pi r^2 L$ provided that L is much smaller than r. $V = 4 \times (3.141) \times (1731000)^2 \times 0.001 = 3.76 \times 10^{10} \text{ m}^3$

Problem 2 – The density of the lunar surface (called the regolith) is about 3000 kilograms/meter³. How many metric tons of regolith are found in the surface volume calculated in Problem 1? Answer: $3.76 \times 10^{10} \text{ m}^3 \times (3000 \text{ kg/m}^3) \times (1 \text{ ton/1000 kg}) = 1.13 \times 10^{11} \text{ metric tons}$.

Problem 3 – The concentration of water is 1 liter per metric ton. How many liters of water could be recovered from the 1 millimeter thick surface layer if 25% of the surface contains water? Answer: 1.13×10^{11} tons x (1 liter water/1 ton regolith) x $1/4 = 2.8 \times 10^{10}$ liters of water.

Problem 4 – How many gallons could be recovered if the entire surface layer were mined? (1 Gallon = 3.78 liters). Answer: 2.8×10^{10} liters x (1 gallon / 3.78 liters) = 7.5×10^{9} gallons of water or about 8 billion gallons of water.

Note: This is similar to the roughly '7 billion gallon' estimate made by the M3 scientists as described in the NASA Press Release for this discovery in September 2009.

For more information, visit:

Moon Minerology Mapper News - http://moonmineralogymapper.jpl.nasa.gov/

The front picture of the moon is from NASA's Moon Mineralogy Mapper on the Indian Space Research Organization's Chandrayaan-1 mission. It is a three-color composite of reflected near-infrared radiation from the sun, and illustrates the extent to which different materials are mapped across the side of the moon that faces Earth. Small amounts of water and hydroxyl (blue) were detected on the surface of the moon at various locations. This image illustrates their distribution at high latitudes toward the poles. Blue shows the signature of water and hydroxyl molecules as seen by a highly diagnostic absorption of infrared light with a wavelength of three micrometers. Green shows the brightness of the surface as measured by reflected infrared radiation from the sun with a wavelength of 2.4 micrometers. Red shows an iron-bearing mineral called pyroxene, detected by absorption of 2.0-micrometer infrared light.